

A Multidimensional Analogue of the Simpson's Formula of Integral

Kazuyuki FUJII *

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

The Simpson's formula is obtained by approximating the integral of a function on some interval by the integral of the quadratic polynomial determined by the function. However, a multidimensional analogue of the formula has not been given as far as we know.

In this paper such a formula is given. Our formula is simple and beautiful, so it may be convenient in Mathematics or Mathematical Physics.


1 Introduction

Let $y = f(x)$ be a continuous function defined on the interval $[a, b]$. We want to calculate the integral $\int_a^b f(x)dx$. However, it is not easy to calculate the integral explicitly, so to make


*E-mail address : fujii@yokohama-cu.ac.jp

some approximation is a realistic way. We recall the Simpson's formula ¹, see for example [1].

We divide $[a, b]$ into equal $2n$ subintervals and set $\Delta = \frac{b-a}{2n}$, and $a = x_0, b = x_{2n}$.



Now we study the integral on small interval $[x_{2k-2}, x_{2k}]$ in the following.



For

$$y_{2k-2} = f(x_{2k-2}), \quad y_{2k-1} = f(x_{2k-1}), \quad y_{2k} = f(x_{2k})$$

we set

$$A = (x_{2k-2}, y_{2k-2}) = (x_{2k-1} - \Delta, y_{2k-2}),$$

$$C = (x_{2k-1}, y_{2k-1}),$$

$$B = (x_{2k}, y_{2k}) = (x_{2k-1} + \Delta, y_{2k})$$

¹The author uses the terminology **Simpson's formula** in place of Simpson's rule because it is nothing but an approximate formula (or method)

for simplicity.

A quadratic polynomial

$$y = px^2 + qx + r$$


passing through three points A, B, C is determined uniquely. Namely, we have only to solve the simultaneous equations

$$\begin{cases} (x_{2k-1} - \Delta)^2 p + (x_{2k-1} - \Delta) q + r = y_{2k-2}, \\ x_{2k-1}^2 p + x_{2k-1} q + r = y_{2k-1}, \\ (x_{2k-1} + \Delta)^2 p + (x_{2k-1} + \Delta) q + r = y_{2k} \end{cases}$$

and the result is

$$\begin{aligned} p &= \frac{y_{2k} - 2y_{2k-1} + y_{2k-2}}{2\Delta^2}, \\ q &= \frac{y_{2k} - y_{2k-2}}{2\Delta} - 2\frac{y_{2k} - 2y_{2k-1} + y_{2k-2}}{2\Delta^2}x_{2k-1}, \\ r &= y_{2k-1} - \frac{y_{2k} - y_{2k-2}}{2\Delta}x_{2k-1} + \frac{y_{2k} - 2y_{2k-1} + y_{2k-2}}{2\Delta^2}x_{2k-1}^2. \end{aligned}$$

Therefore the function $y = f(x)$ on the interval $[x_{2k-2}, x_{2k}]$ is approximated by $y = px^2 + qx + r$ like



Then an important formula is obtained

$$\int_{x_{2k-1}-\Delta}^{x_{2k-1}+\Delta} (px^2 + qx + r) dx = \frac{\Delta}{3} (y_{2k-2} + 4y_{2k-1} + y_{2k})$$

because

$$\begin{aligned}
\text{LHS} &= \frac{p}{3} \{(x_{2k-1} + \Delta)^3 - (x_{2k-1} - \Delta)^3\} + \frac{q}{2} \{(x_{2k-1} + \Delta)^2 - (x_{2k-1} - \Delta)^2\} + 2r\Delta \\
&= \frac{2}{3}p\Delta^3 + 2(px_{2k-1}^2 + qx_{2k-1} + r)\Delta \\
&= \frac{2}{3}p\Delta^3 + 2y_{2k-1}\Delta \\
&= \frac{\Delta}{3} \left\{ 2\Delta^2 \times \frac{y_{2k} - 2y_{2k-1} + y_{2k-2}}{2\Delta^2} + 6y_{2k-1} \right\} \\
&= \frac{\Delta}{3} (y_{2k-2} + 4y_{2k-1} + y_{2k}) = \text{RHS}.
\end{aligned}$$

From this we have the famous Simpson's formula

$$\begin{aligned}
\int_a^b f(x)dx &= \sum_{k=1}^n \int_{x_{2k-2}}^{x_{2k}} f(x)dx = \sum_{k=1}^n \int_{x_{2k-1}-\Delta}^{x_{2k-1}+\Delta} f(x)dx \\
&\approx \sum_{k=1}^n \int_{x_{2k-1}-\Delta}^{x_{2k-1}+\Delta} (px^2 + qx + r) dx \\
&= \sum_{k=1}^n \frac{\Delta}{3} (y_{2k-2} + 4y_{2k-1} + y_{2k}) \\
&= \frac{b-a}{6n} \{y_0 + 4(y_1 + y_3 + \cdots + y_{2n-1}) + 2(y_2 + y_4 + \cdots + y_{2n-2}) + y_{2n}\}
\end{aligned}$$

because $\Delta = \frac{b-a}{2n}$.

2 Generalized Simpson's Formula

Let D be an n -dimensional cuboid

$$D = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n] \subset \mathbf{R}^n$$


and f be a continuous function on D . What we do is to calculate the multidimensional integral

$$\int \int \cdots \int_D f(x_1, x_2, \cdots, x_n) dx_1 dx_2 \cdots dx_n. \quad (1)$$

However, it is almost difficult, so we must satisfy only by approximating it as shown in the introduction. For that purpose we rewrite D as follows.

$$D = [\alpha_1 - \Delta_1, \alpha_1 + \Delta_1] \times [\alpha_2 - \Delta_2, \alpha_2 + \Delta_2] \times \cdots \times [\alpha_n - \Delta_n, \alpha_n + \Delta_n] \quad (2)$$

where $\alpha_j = (b_j + a_j)/2$ and $\Delta_j = (b_j - a_j)/2$. See the following figure ($n = 2$).



Next we approximate f by a quadratic polynomial

$$F(x_1, x_2, \dots, x_n) = \sum_{i_1, i_2, \dots, i_n=0}^2 a_{i_1 i_2 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}, \quad (3)$$

which is a natural extension in case of $n = 1$.

For $j_1, j_2, \dots, j_n \in \{-1, 0, 1\}$ we set

$$\begin{aligned} w_{j_1 j_2 \dots j_n} &= f(\alpha_1 + j_1 \Delta_1, \alpha_2 + j_2 \Delta_2, \dots, \alpha_n + j_n \Delta_n) \\ &= F(\alpha_1 + j_1 \Delta_1, \alpha_2 + j_2 \Delta_2, \dots, \alpha_n + j_n \Delta_n). \end{aligned} \quad (4)$$

It is of course

$$\#\{w_{j_1 j_2 \dots j_n}\} = 3^n = \#\{a_{i_1 i_2 \dots i_n}\},$$

see the figure above. Then we have

Formula (Conjecture)

$$\begin{aligned} &\int \int \dots \int_D F(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n \\ &= \frac{\Delta_1 \Delta_2 \dots \Delta_n}{3^n} \sum_{j_1, j_2, \dots, j_n \in \{-1, 0, 1\}} 4^{\#\{j_1, j_2, \dots, j_n\}_0} w_{j_1 j_2 \dots j_n} \end{aligned} \quad (5)$$

where $\#\{j_1, j_2, \dots, j_n\}_0$ is the number of 0 in $\{j_1, j_2, \dots, j_n\}$.

For the case of $n = 2$ and $n = 3$ the formula is proved in the following. However, the proof of the general case is left to readers.

From the formula we obtain a good approximation

$$\begin{aligned} & \int \int \cdots \int_D f(x_1, x_2, \cdots, x_n) dx_1 dx_2 \cdots dx_n \\ & \approx \frac{\Delta_1 \Delta_2 \cdots \Delta_n}{3^n} \sum_{j_1, j_2, \cdots, j_n \in \{-1, 0, 1\}} 4^{\#\{j_1, j_2, \cdots, j_n\}_0} w_{j_1 j_2 \cdots j_n} \end{aligned} \quad (6)$$


if D is small enough.

2.1 Proof for $n = 2$

Let us prove (5) for $n = 2$. From (3) we set for simplicity

$$\begin{aligned} w &= F(x, y) = \sum_{i, j=0}^2 a_{ij} x^i y^j \\ &= a_{20} x^2 + a_{10} x + a_{00} + (a_{21} x^2 + a_{11} x + a_{01}) y + (a_{22} x^2 + a_{12} x + a_{02}) y^2. \end{aligned} \quad (7)$$

See the figure once more.



Then

$$\begin{aligned}
& \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} w dx dy \\
&= \left\{ a_{20} \frac{(\alpha_1 + \Delta_1)^3 - (\alpha_1 - \Delta_1)^3}{3} + a_{10} \frac{(\alpha_1 + \Delta_1)^2 - (\alpha_1 - \Delta_1)^2}{2} + a_{00} \cdot 2\Delta_1 \right\} 2\Delta_2 \\
&+ \left\{ a_{21} \frac{(\alpha_1 + \Delta_1)^3 - (\alpha_1 - \Delta_1)^3}{3} + a_{11} \frac{(\alpha_1 + \Delta_1)^2 - (\alpha_1 - \Delta_1)^2}{2} + a_{01} \cdot 2\Delta_1 \right\} \times \\
&\quad \frac{(\alpha_2 + \Delta_2)^2 - (\alpha_2 - \Delta_2)^2}{2} \\
&+ \left\{ a_{22} \frac{(\alpha_1 + \Delta_1)^3 - (\alpha_1 - \Delta_1)^3}{3} + a_{12} \frac{(\alpha_1 + \Delta_1)^2 - (\alpha_1 - \Delta_1)^2}{2} + a_{02} \cdot 2\Delta_1 \right\} \times \\
&\quad \frac{(\alpha_2 + \Delta_2)^3 - (\alpha_2 - \Delta_2)^3}{3} \\
&= \dots \\
&= 4\Delta_1\Delta_2 \left\{ a_{20}\alpha_1^2 + a_{10}\alpha_1 + a_{00} + (a_{21}\alpha_1^2 + a_{11}\alpha_1 + a_{01})\alpha_2 + (a_{22}\alpha_1^2 + a_{12}\alpha_1 + a_{02})\alpha_2^2 \right\} \\
&+ \frac{4\Delta_1\Delta_2}{3} (a_{20}\Delta_1^2 + a_{21}\Delta_1^2\alpha_2 + a_{22}\Delta_1^2\alpha_2^2 + a_{02}\Delta_2^2 + a_{12}\alpha_1\Delta_2^2 + a_{22}\alpha_1^2\Delta_2^2) \\
&+ \frac{4\Delta_1\Delta_2}{9} a_{22}\Delta_1^2\Delta_2^2. \tag{8}
\end{aligned}$$

From (4) we have 9(= 3²)-data

$$w_{k,l} = w(\alpha_1 + k\Delta_1, \alpha_2 + l\Delta_2) \tag{9}$$

for $k, l \in \{-1, 0, 1\}$, so we want to express the integral above in terms of these data $\{w_{kl}\}$.

Since

$$a_{20}\alpha_1^2 + a_{10}\alpha_1 + a_{00} + (a_{21}\alpha_1^2 + a_{11}\alpha_1 + a_{01})\alpha_2 + (a_{22}\alpha_1^2 + a_{12}\alpha_1 + a_{02})\alpha_2^2 = w_{0,0} \tag{10}$$

is clear we must treat the remaining ones

$$\begin{aligned}
& a_{20}\Delta_1^2 + a_{21}\Delta_1^2\alpha_2 + a_{22}\Delta_1^2\alpha_2^2 + a_{02}\Delta_2^2 + a_{12}\alpha_1\Delta_2^2 + a_{22}\alpha_1^2\Delta_2^2, \\
& a_{22}\Delta_1^2\Delta_2^2.
\end{aligned}$$

It is not difficult to show

$$\begin{aligned}
a_{20}\Delta_1^2 + a_{21}\Delta_1^2\alpha_2 + a_{22}\Delta_1^2\alpha_2^2 &= \frac{w_{-1,0} - 2w_{0,0} + w_{1,0}}{2}, \\
a_{02}\Delta_2^2 + a_{12}\alpha_1\Delta_2^2 + a_{22}\alpha_1^2\Delta_2^2 &= \frac{w_{0,-1} - 2w_{0,0} + w_{0,1}}{2},
\end{aligned}$$

so we have

$$\begin{aligned}
& a_{20}\Delta_1^2 + a_{21}\Delta_1^2\alpha_2 + a_{22}\Delta_1^2\alpha_2^2 + a_{02}\Delta_2^2 + a_{12}\alpha_1\Delta_2^2 + a_{22}\alpha_1^2\Delta_2^2 \\
&= \frac{w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1} - 4w_{0,0}}{2}.
\end{aligned} \tag{11}$$

Similarly,

$$\begin{aligned}
& \frac{w_{-1,-1} + w_{-1,1} + w_{1,-1} + w_{1,1}}{4} \\
&= a_{20}\alpha_1^2 + a_{10}\alpha_1 + a_{00} + (a_{21}\alpha_1^2 + a_{11}\alpha_1 + a_{01})\alpha_2 + (a_{22}\alpha_1^2 + a_{12}\alpha_1 + a_{02})\alpha_2^2 \\
&+ a_{20}\Delta_1^2 + a_{21}\Delta_1^2\alpha_2 + a_{22}\Delta_1^2\alpha_2^2 + a_{02}\Delta_2^2 + a_{12}\alpha_1\Delta_2^2 + a_{22}\alpha_1^2\Delta_2^2 \\
&+ a_{22}\Delta_1^2\Delta_2^2 \\
&= w_{0,0} + \frac{w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1} - 4w_{0,0}}{2} + a_{22}\Delta_1^2\Delta_2^2
\end{aligned}$$

so we have

$$a_{22}\Delta_1^2\Delta_2^2 = \frac{w_{-1,-1} + w_{-1,1} + w_{1,-1} + w_{1,1} - 2(w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1}) + 4w_{0,0}}{4}. \tag{12}$$

Therefore, substituting (10), (11) and (12) into (8) we obtain

$$\begin{aligned}
& \int_{\alpha_1-\Delta_1}^{\alpha_1+\Delta_1} \int_{\alpha_2-\Delta_2}^{\alpha_2+\Delta_2} w dx dy \\
&= 4\Delta_1\Delta_2 w_{0,0} + \frac{2\Delta_1\Delta_2}{3}(w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1} - 4w_{0,0}) \\
&+ \frac{\Delta_1\Delta_2}{9}(w_{-1,-1} + w_{-1,1} + w_{1,-1} + w_{1,1} - 2(w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1}) + 4w_{0,0}) \\
&= \frac{\Delta_1\Delta_2}{9} \{w_{-1,-1} + w_{-1,1} + w_{1,-1} + w_{1,1} + 4(w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1}) + 16w_{0,0}\}.
\end{aligned} \tag{13}$$

This formula is interesting enough.

2.2 Proof for $n = 3$

Let us prove (5) for $n = 3$. From (3) we set

$$\begin{aligned}
w &= F(x, y, z) = \sum_{i,j,k=0}^2 a_{ijk} x^i y^j z^k \\
&= a_{200}x^2 + a_{100}x + a_{000} + (a_{210}x^2 + a_{110}x + a_{010})y + (a_{220}x^2 + a_{120}x + a_{020})y^2 \\
&+ \{a_{201}x^2 + a_{101}x + a_{001} + (a_{211}x^2 + a_{111}x + a_{011})y + (a_{221}x^2 + a_{121}x + a_{021})y^2\} z \\
&+ \{a_{202}x^2 + a_{102}x + a_{002} + (a_{212}x^2 + a_{112}x + a_{012})y + (a_{222}x^2 + a_{122}x + a_{022})y^2\} z^2.
\end{aligned} \tag{14}$$

Then

$$\begin{aligned}
&\int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \int_{\alpha_3 - \Delta_3}^{\alpha_3 + \Delta_3} w dx dy dz \\
&= \left\{ \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \sum_{i,j=0}^2 a_{ij0} x^i y^j \right\} 2\Delta_3 \\
&+ \left\{ \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \sum_{i,j=0}^2 a_{ij1} x^i y^j \right\} \frac{(\alpha_3 + \Delta_3)^2 - (\alpha_3 - \Delta_3)^2}{2} \\
&+ \left\{ \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \sum_{i,j=0}^2 a_{ij2} x^i y^j \right\} \frac{(\alpha_3 + \Delta_3)^3 - (\alpha_3 - \Delta_3)^3}{3}.
\end{aligned}$$

By (8) some calculation gives

$$\begin{aligned}
& \int_{\alpha_1-\Delta_1}^{\alpha_1+\Delta_1} \int_{\alpha_2-\Delta_2}^{\alpha_2+\Delta_2} \int_{\alpha_3-\Delta_3}^{\alpha_3+\Delta_3} w dx dy dz \\
= & [4\Delta_1\Delta_2\{a_{200}\alpha_1^2 + a_{100}\alpha_1 + a_{000} + (a_{210}\alpha_1^2 + a_{110}\alpha_1 + a_{010})\alpha_2 + (a_{220}\alpha_1^2 + a_{120}\alpha_1 + a_{020})\alpha_2^2\} \\
& + \frac{4\Delta_1\Delta_2}{3}(a_{200}\Delta_1^2 + a_{210}\Delta_1^2\alpha_2 + a_{220}\Delta_1^2\alpha_2^2 + a_{020}\Delta_2^2 + a_{120}\alpha_1\Delta_2^2 + a_{220}\alpha_1^2\Delta_2^2) \\
& + \frac{4\Delta_1\Delta_2}{9}a_{220}\Delta_1^2\Delta_2^2]2\Delta_3 \\
& + [4\Delta_1\Delta_2\{a_{201}\alpha_1^2 + a_{101}\alpha_1 + a_{001} + (a_{211}\alpha_1^2 + a_{111}\alpha_1 + a_{011})\alpha_2 + (a_{221}\alpha_1^2 + a_{121}\alpha_1 + a_{021})\alpha_2^2\} \\
& + \frac{4\Delta_1\Delta_2}{3}(a_{201}\Delta_1^2 + a_{211}\Delta_1^2\alpha_2 + a_{221}\Delta_1^2\alpha_2^2 + a_{021}\Delta_2^2 + a_{121}\alpha_1\Delta_2^2 + a_{221}\alpha_1^2\Delta_2^2) \\
& + \frac{4\Delta_1\Delta_2}{9}a_{221}\Delta_1^2\Delta_2^2]2\alpha_3\Delta_3 \\
& + [4\Delta_1\Delta_2\{a_{202}\alpha_1^2 + a_{102}\alpha_1 + a_{002} + (a_{212}\alpha_1^2 + a_{112}\alpha_1 + a_{012})\alpha_2 + (a_{222}\alpha_1^2 + a_{122}\alpha_1 + a_{022})\alpha_2^2\} \\
& + \frac{4\Delta_1\Delta_2}{3}(a_{202}\Delta_1^2 + a_{212}\Delta_1^2\alpha_2 + a_{222}\Delta_1^2\alpha_2^2 + a_{022}\Delta_2^2 + a_{122}\alpha_1\Delta_2^2 + a_{222}\alpha_1^2\Delta_2^2) \\
& + \frac{4\Delta_1\Delta_2}{9}a_{222}\Delta_1^2\Delta_2^2](2\alpha_3^2\Delta_3 + \frac{2}{3}\Delta_3^3) \\
= & 8\Delta_1\Delta_2\Delta_3 \times \\
& [\{a_{200}\alpha_1^2 + a_{100}\alpha_1 + a_{000} + (a_{210}\alpha_1^2 + a_{110}\alpha_1 + a_{010})\alpha_2 + (a_{220}\alpha_1^2 + a_{120}\alpha_1 + a_{020})\alpha_2^2\} \\
& + \{a_{201}\alpha_1^2 + a_{101}\alpha_1 + a_{001} + (a_{211}\alpha_1^2 + a_{111}\alpha_1 + a_{011})\alpha_2 + (a_{221}\alpha_1^2 + a_{121}\alpha_1 + a_{021})\alpha_2^2\}\alpha_3 \\
& + \{a_{202}\alpha_1^2 + a_{102}\alpha_1 + a_{002} + (a_{212}\alpha_1^2 + a_{112}\alpha_1 + a_{012})\alpha_2 + (a_{222}\alpha_1^2 + a_{122}\alpha_1 + a_{022})\alpha_2^2\}\alpha_3^2] \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{3} \times \\
& [\{a_{200}\Delta_1^2 + a_{210}\Delta_1^2\alpha_2 + a_{220}\Delta_1^2\alpha_2^2 + a_{020}\Delta_2^2 + a_{120}\alpha_1\Delta_2^2 + a_{220}\alpha_1^2\Delta_2^2\} \\
& + \{a_{201}\Delta_1^2 + a_{211}\Delta_1^2\alpha_2 + a_{221}\Delta_1^2\alpha_2^2 + a_{021}\Delta_2^2 + a_{121}\alpha_1\Delta_2^2 + a_{221}\alpha_1^2\Delta_2^2\}\alpha_3 \\
& + \{a_{202}\Delta_1^2 + a_{212}\Delta_1^2\alpha_2 + a_{222}\Delta_1^2\alpha_2^2 + a_{022}\Delta_2^2 + a_{122}\alpha_1\Delta_2^2 + a_{222}\alpha_1^2\Delta_2^2\}\alpha_3^2] \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{9}(a_{220}\Delta_1^2\Delta_2^2 + a_{221}\Delta_1^2\Delta_2^2\alpha_3 + a_{222}\Delta_1^2\Delta_2^2\alpha_3^2) \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{3} \times \\
& \{a_{202}\alpha_1^2 + a_{102}\alpha_1 + a_{002} + (a_{212}\alpha_1^2 + a_{112}\alpha_1 + a_{012})\alpha_2 + (a_{222}\alpha_1^2 + a_{122}\alpha_1 + a_{022})\alpha_2^2\}\Delta_3^2 \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{9}(a_{202}\Delta_1^2 + a_{212}\Delta_1^2\alpha_2 + a_{222}\Delta_1^2\alpha_2^2 + a_{022}\Delta_2^2 + a_{122}\alpha_1\Delta_2^2 + a_{222}\alpha_1^2\Delta_2^2)\Delta_3^2 \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{27}a_{222}\Delta_1^2\Delta_2^2\Delta_3^2.
\end{aligned}$$

By rearranging terms we have

$$\begin{aligned}
& \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \int_{\alpha_3 - \Delta_3}^{\alpha_3 + \Delta_3} w dx dy dz \\
= & 8\Delta_1\Delta_2\Delta_3 \times \\
& [\{ a_{200}\alpha_1^2 + a_{100}\alpha_1 + a_{000} + (a_{210}\alpha_1^2 + a_{110}\alpha_1 + a_{010})\alpha_2 + (a_{220}\alpha_1^2 + a_{120}\alpha_1 + a_{020})\alpha_2^2 \} \\
& + \{ a_{201}\alpha_1^2 + a_{101}\alpha_1 + a_{001} + (a_{211}\alpha_1^2 + a_{111}\alpha_1 + a_{011})\alpha_2 + (a_{221}\alpha_1^2 + a_{121}\alpha_1 + a_{021})\alpha_2^2 \} \alpha_3 \\
& + \{ a_{202}\alpha_1^2 + a_{102}\alpha_1 + a_{002} + (a_{212}\alpha_1^2 + a_{112}\alpha_1 + a_{012})\alpha_2 + (a_{222}\alpha_1^2 + a_{122}\alpha_1 + a_{022})\alpha_2^2 \} \alpha_3^2] \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{3} \times \\
& [\{ a_{200} + a_{210}\alpha_2 + a_{220}\alpha_2^2 + (a_{201} + a_{211}\alpha_2 + a_{221}\alpha_2^2)\alpha_3 + (a_{202} + a_{212}\alpha_2 + a_{222}\alpha_2^2)\alpha_3^2 \} \Delta_1^2 \\
& + \{ a_{020} + a_{120}\alpha_1 + a_{220}\alpha_1^2 + (a_{021} + a_{121}\alpha_1 + a_{221}\alpha_1^2)\alpha_3 + (a_{022} + a_{122}\alpha_1 + a_{222}\alpha_1^2)\alpha_3^2 \} \Delta_2^2 \\
& + \{ a_{002} + a_{102}\alpha_1 + a_{202}\alpha_1^2 + (a_{012} + a_{112}\alpha_1 + a_{212}\alpha_1^2)\alpha_2 + (a_{022} + a_{122}\alpha_1 + a_{222}\alpha_1^2)\alpha_2^2 \} \Delta_3^2] \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{9} \times \\
& \{ a_{220}\Delta_1^2\Delta_2^2 + a_{221}\Delta_1^2\Delta_2^2\alpha_3 + a_{222}\Delta_1^2\Delta_2^2\alpha_3^2 + a_{202}\Delta_1^2\Delta_3^2 + a_{212}\Delta_1^2\alpha_2\Delta_3^2 + a_{222}\Delta_1^2\alpha_2^2\Delta_3^2 \\
& + a_{022}\Delta_2^2\Delta_3^2 + a_{122}\alpha_1\Delta_2^2\Delta_3^2 + a_{222}\alpha_1^2\Delta_2^2\Delta_3^2 \} \\
& + \frac{8\Delta_1\Delta_2\Delta_3}{27} a_{222}\Delta_1^2\Delta_2^2\Delta_3^2. \tag{15}
\end{aligned}$$

From (4) we have $27(=3^3)$ -data

$$w_{l,m,n} = w(\alpha_1 + l\Delta_1, \alpha_2 + m\Delta_2, \alpha_3 + n\Delta_3) \tag{16}$$

for $l, m, n \in \{-1, 0, 1\}$, so we must express the integral above in terms of these data $\{w_{l,m,n}\}$.

Since

$$\begin{aligned}
& \{ a_{200}\alpha_1^2 + a_{100}\alpha_1 + a_{000} + (a_{210}\alpha_1^2 + a_{110}\alpha_1 + a_{010})\alpha_2 + (a_{220}\alpha_1^2 + a_{120}\alpha_1 + a_{020})\alpha_2^2 \} \\
& + \{ a_{201}\alpha_1^2 + a_{101}\alpha_1 + a_{001} + (a_{211}\alpha_1^2 + a_{111}\alpha_1 + a_{011})\alpha_2 + (a_{221}\alpha_1^2 + a_{121}\alpha_1 + a_{021})\alpha_2^2 \} \alpha_3 \\
& + \{ a_{202}\alpha_1^2 + a_{102}\alpha_1 + a_{002} + (a_{212}\alpha_1^2 + a_{112}\alpha_1 + a_{012})\alpha_2 + (a_{222}\alpha_1^2 + a_{122}\alpha_1 + a_{022})\alpha_2^2 \} \alpha_3^2 \\
= & w_{0,0,0} \tag{17}
\end{aligned}$$

we treat the remaining ones. Here we list formulas necessary to prove the main formula.

$$\begin{aligned}
& \{a_{200} + a_{210}\alpha_2 + a_{220}\alpha_2^2 + (a_{201} + a_{211}\alpha_2 + a_{221}\alpha_2^2)\alpha_3 + (a_{202} + a_{212}\alpha_2 + a_{222}\alpha_2^2)\alpha_3^2\}\Delta_1^2 \\
&= \frac{1}{2}(w_{-1,0,0} + w_{1,0,0}) - w_{0,0,0}, \\
& \{a_{020} + a_{120}\alpha_1 + a_{220}\alpha_1^2 + (a_{021} + a_{121}\alpha_1 + a_{221}\alpha_1^2)\alpha_3 + (a_{022} + a_{122}\alpha_1 + a_{222}\alpha_1^2)\alpha_3^2\}\Delta_2^2 \\
&= \frac{1}{2}(w_{0,-1,0} + w_{0,1,0}) - w_{0,0,0}, \\
& \{a_{002} + a_{102}\alpha_1 + a_{202}\alpha_1^2 + (a_{012} + a_{112}\alpha_1 + a_{212}\alpha_1^2)\alpha_2 + (a_{022} + a_{122}\alpha_1 + a_{222}\alpha_1^2)\alpha_2^2\}\Delta_3^2 \\
&= \frac{1}{2}(w_{0,0,-1} + w_{0,0,1}) - w_{0,0,0} \tag{18}
\end{aligned}$$

and

$$\begin{aligned}
& a_{220}\Delta_1^2\Delta_2^2 + a_{221}\Delta_1^2\Delta_2^2\alpha_3 + a_{222}\Delta_1^2\Delta_2^2\alpha_3^2 + a_{202}\Delta_1^2\Delta_3^2 + a_{212}\Delta_1^2\alpha_2\Delta_3^2 + a_{222}\Delta_1^2\alpha_2^2\Delta_3^2 \\
& + a_{022}\Delta_2^2\Delta_3^2 + a_{122}\alpha_1\Delta_2^2\Delta_3^2 + a_{222}\alpha_1^2\Delta_2^2\Delta_3^2 \\
&= \frac{1}{4}(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \\
& \quad + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \\
& \quad - (w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) + 3w_{0,0,0} \tag{19}
\end{aligned}$$

and

$$\begin{aligned}
& a_{222}\Delta_1^2\Delta_2^2\Delta_3^2 \\
&= \frac{1}{8}(w_{-1,-1,-1} + w_{1,-1,-1} + w_{1,1,-1} + w_{-1,1,-1} + w_{-1,-1,1} + w_{1,-1,1} + w_{1,1,1} + w_{-1,1,1}) \\
& \quad - \frac{1}{4}(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \\
& \quad + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \\
& \quad + \frac{1}{2}(w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) - w_{0,0,0}. \tag{20}
\end{aligned}$$

The proofs are long but straightforward, so they are left to readers.

Therefore, substituting (17) $\sim$ (20) into (15) we obtain

$$\begin{aligned}
& \int_{\alpha_1-\Delta_1}^{\alpha_1+\Delta_1} \int_{\alpha_2-\Delta_2}^{\alpha_2+\Delta_2} \int_{\alpha_3-\Delta_3}^{\alpha_3+\Delta_3} w dx dy dz \\
= & 8\Delta_1\Delta_2\Delta_3 w_{0,0,0} \\
+ & \frac{8\Delta_1\Delta_2\Delta_3}{3} \left\{ \frac{1}{2}(w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) - 3w_{0,0,0} \right\} \\
+ & \frac{8\Delta_1\Delta_2\Delta_3}{9} \left\{ \frac{1}{4}(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \right. \\
& \quad \left. + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \right. \\
& \quad \left. - (w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) + 3w_{0,0,0} \right\} \\
+ & \frac{8\Delta_1\Delta_2\Delta_3}{27} \left\{ \frac{1}{8}(w_{-1,-1,-1} + w_{1,-1,-1} + w_{1,1,-1} + w_{-1,1,-1} + w_{-1,-1,1} + w_{1,-1,1} + w_{1,1,1} + w_{-1,1,1}) \right. \\
& \quad \left. - \frac{1}{4}(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \right. \\
& \quad \left. + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \right. \\
& \quad \left. + \frac{1}{2}(w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) - w_{0,0,0} \right\} \\
= & \frac{\Delta_1\Delta_2\Delta_3}{27} \times \\
& \{ w_{-1,-1,-1} + w_{1,-1,-1} + w_{1,1,-1} + w_{-1,1,-1} + w_{-1,-1,1} + w_{1,-1,1} + w_{1,1,1} + w_{-1,1,1} \\
& + 4(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \\
& \quad + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \\
& + 16(w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) \\
& + 64w_{0,0,0} \}. \tag{21}
\end{aligned}$$

The proof is a bit complicated.

2.3 General Case

Let us consider the general case. Unfortunately, we have no method to calculate the general case at the present time. However, from the results (13) and (21) we can conjecture the general formula as follows.

The formulas (13) and (21) can be rewritten as

$$\begin{aligned} & \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} w dx dy \\ &= \frac{\Delta_1 \Delta_2}{3^2} \{ w_{-1,-1} + w_{-1,1} + w_{1,-1} + w_{1,1} + 4(w_{-1,0} + w_{1,0} + w_{0,-1} + w_{0,1}) + 4^2 w_{0,0} \} \end{aligned}$$

and

$$\begin{aligned} & \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \int_{\alpha_3 - \Delta_3}^{\alpha_3 + \Delta_3} w dx dy dz \\ &= \frac{\Delta_1 \Delta_2 \Delta_3}{3^3} \times \\ & \{ w_{-1,-1,-1} + w_{1,-1,-1} + w_{1,1,-1} + w_{-1,1,-1} + w_{-1,-1,1} + w_{1,-1,1} + w_{1,1,1} + w_{-1,1,1} \\ & + 4(w_{-1,-1,0} + w_{-1,1,0} + w_{1,-1,0} + w_{1,1,0} + w_{-1,0,-1} + w_{-1,0,1} + w_{1,0,-1} + w_{1,0,1} \\ & + w_{0,-1,-1} + w_{0,-1,1} + w_{0,1,-1} + w_{0,1,1}) \\ & + 4^2(w_{-1,0,0} + w_{1,0,0} + w_{0,-1,0} + w_{0,1,0} + w_{0,0,-1} + w_{0,0,1}) \\ & + 4^3 w_{0,0,0} \}. \end{aligned}$$

From the forms it is easy to conjecture the general form.

If we set

$$w = F(x_1, x_2, \dots, x_n) = \sum_{i_1, i_2, \dots, i_n=0}^2 a_{i_1 i_2 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \quad (22)$$

then

$$\begin{aligned} & \int_{\alpha_1 - \Delta_1}^{\alpha_1 + \Delta_1} \int_{\alpha_2 - \Delta_2}^{\alpha_2 + \Delta_2} \dots \int_{\alpha_n - \Delta_n}^{\alpha_n + \Delta_n} w dx_1 dx_2 \dots dx_n \\ &= \frac{\Delta_1 \Delta_2 \dots \Delta_n}{3^n} \sum_{j_1, j_2, \dots, j_n \in \{-1, 0, 1\}} 4^{\#\{j_1, j_2, \dots, j_n\}_0} w_{j_1 j_2 \dots j_n} \end{aligned} \quad (23)$$

where $\#\{j_1, j_2, \dots, j_n\}_0$ is the number of 0 in $\{j_1, j_2, \dots, j_n\}$ and

$$\begin{aligned} w_{j_1 j_2 \dots j_n} &= F(\alpha_1 + j_1 \Delta_1, \alpha_2 + j_2 \Delta_2, \dots, \alpha_n + j_n \Delta_n) \\ &= f(\alpha_1 + j_1 \Delta_1, \alpha_2 + j_2 \Delta_2, \dots, \alpha_n + j_n \Delta_n). \end{aligned}$$

The formula is beautiful enough.

3 Concluding Remarks

In the paper we gave a multidimensional analogue of the Simpson's formula of integral. As far as we know the formula has not been given, which is a bit mysterious. We believe that it will become useful in all fields related to integrals.

Next let us consider more general problems. Let (M, g) be a curved space-time and f be a function on M . If we consider the integral

$$\int_M f(x) dv_g(x)$$

where dv_g is some measure on M , it is almost impossible to calculate. For that we want to discretize the integral as follows.

$$\sum_{\{D\}} \int_D f(x) dv_g(x) \longrightarrow \sum_{\{D\}} \{\text{some formula like (23)}\}$$

Therefore we must choose D to be “computable”, which is just key point.

Details of calculation will be reported in [2].

References

- [1] WolframMathWorld : <http://mathworld.wolfram.com/>
- [2] K. Fujii : in progress.